

Separation of proton polarizabilities with the beam asymmetry of Compton scattering

Nadiia Krupina and Vladimir Pascalutsa

*Excellence Cluster PRISMA Institut für Kernphysik,
Johannes Gutenberg-Universität Mainz, 55128 Mainz, Germany*

(Dated: April 30, 2013)

We propose to determine the magnetic dipole polarizability of the proton directly from the beam asymmetry of low-energy Compton scattering based on the fact that the leading non-Born contribution to the asymmetry is due to the magnetic polarizability alone; the electric polarizability cancels out. The beam asymmetry thus provides the simplest observable with a clean separation of the magnetic polarizability from the electric one. Introducing polarizabilities in a Lorentz-invariant fashion we compute the higher-order (recoil) effects of polarizabilities on beam asymmetry and show that these effects are most suppressed in forward kinematics. With the prospects of precision Compton experiments at the MAMI and HIGS facilities in mind, we argue why the beam asymmetry could be the best way to measure the elusive magnetic polarizability of the proton.

PACS numbers: 13.60.Fz - Elastic and Compton scattering, 14.20.Dh - Protons and neutrons, 25.20.Dc - Photon absorption and scattering

The current Particle Data Group (PDG) [1] values of the electric- and magnetic-dipole polarizabilities of the proton [2, 3], i.e.,

$$\alpha_{E1} = (12.0 \pm 0.6) \times 10^{-4} \text{ fm}^3, \quad (1a)$$

$$\beta_{M1} = (1.9 \pm 0.5) \times 10^{-4} \text{ fm}^3 \quad (1b)$$

are in significant disagreement with the most recent post-dictions of chiral effective field theory (ChEFT) [4, 5], as can be seen in Fig 1. The state-of-the-art ChEFT calculations, based on either the baryon (B) or heavy-baryon (HB) chiral perturbation theory (ChPT) with octet and decuplet fields [11], are in excellent agreement with the experimental Compton-scattering cross sections, but not necessarily in agreement with the polarizabilities extracted from these data by the experimental groups. See [12] for a recent review.

A likely source of these discrepancies is an underestimate of *model dependence* in the extraction of polarizabilities from Compton-scattering data. In principle, one should opt for a model-independent extraction, based on the low-energy expansion (LEX) of Compton-scattering observables, where the leading order terms, beyond the Born term, are expressed through polarizabilities. For example, the non-Born (NB) part of the unpolarized differential cross-section for Compton scattering off a target with mass M and charge Ze is given by

$$\frac{d\sigma^{(\text{NB})}}{d\Omega_L} = -\frac{Z^2\alpha_{em}}{M} \left(\frac{\nu'}{\nu}\right)^2 \nu\nu' [\alpha_{E1} (1 + \cos^2 \theta_L) + 2\beta_{M1} \cos \theta_L] + O(\nu^4), \quad (2)$$

where $\nu = (s - M^2)/2M$ and $\nu' = (-u + M^2)/2M$ are respectively the energies of the incident and scattered photon in the lab frame, θ_L ($d\Omega_L = 2\pi \sin \theta_L d\theta_L$) the scattering (solid) angle; s , u and $t = 2M(\nu' - \nu)$ are the Mandelstam variables; and $\alpha_{em} = e^2/4\pi$ is the fine-structure constant. Hence, given the exactly known Born

contribution [13], and the experimental angular distribution at very low energy one could in principle extract the polarizabilities with a negligible model dependence [2]. In reality however, in order to resolve the small polarizability effect in the tiny Compton cross sections, most of the measurements are done at energies exceeding 100 MeV, i.e., not small compared to the pion mass m_π . It is m_π , the onset of the pion-production branch cut, that severely limits the applicability of a polynomial expansion




FIG. 1: (Color online). The scalar polarizabilities of the proton. Magenta blob represents the PDG summary [1]. Experimental results are from Federspiel et al. [6], Zieger et al. [7], MacGibbon et al. [8], and TAPS [9]. ‘Sum Rule’ indicates the Baldin sum rule evaluations of $\alpha_{E1} + \beta_{M1}$ [9] (broader band) and [10]. ChPT calculations are from [4] (B χ PT—red blob) and [5] (HB χ PT—cyan blob).

sion in energy such as LEX. At the energies around the pion-production threshold one obtains a very substantial sensitivity to polarizabilities but needs to resort to a model-dependent approach in order to extract them. See [14, 15] for reviews.

The magnetic polarizability β_{M1} seems to be affected the most: the central value of the BChPT calculation is a factor of 2 larger than the PDG value. This is attributed to the dominance of α_{E1} in the unpolarized cross section.¹ It is desirable to find an observable sensitive to β_{M1} alone, such that the latter could be determined independently of α_{E1} . According to the leading-order (LO) LEX for cross sections involving linearly polarized photons [18], the difference of cross sections for photons polarized perpendicular or parallel to the scattering plane,

$$(d\sigma_{\perp} - d\sigma_{\parallel})/d\Omega \quad (3)$$

depends only on α_{E1} , while the combination

$$(\cos^2 \theta d\sigma_{\perp} - d\sigma_{\parallel})/d\Omega \quad (4)$$

only on β_{M1} . New experiments at the Mainz Microtron (MAMI) and the High Intensity Gamma Source (HIGS) are planned to measure these two combinations in order to extract α_{E1} and β_{M1} independently. This Letter aims to show that β_{M1} can directly be extracted from the beam asymmetry:

$$\Sigma_3 \equiv \frac{d\sigma_{\parallel} - d\sigma_{\perp}}{d\sigma_{\parallel} + d\sigma_{\perp}}, \quad (5)$$

and that such extraction is potentially more accurate than the one based on Eq. (4).

Indeed, applying the LEX for the beam asymmetry of proton Compton scattering we arrive at the following result:

$$\Sigma_3 = \Sigma_3^{(B)} - \frac{4M\omega^2 \cos \theta \sin^2 \theta}{\alpha_{em}(1 + \cos^2 \theta)^2} \beta_{M1} + O(\omega^4), \quad (6)$$

where $\Sigma_3^{(B)}$ is the pure Born contribution, while

$$\omega = \frac{s - M^2 + \frac{1}{2}t}{\sqrt{4M^2 - t}}, \quad \theta = \arccos \left(1 + \frac{t}{2\omega^2} \right) \quad (7)$$

are the photon energy and scattering angle in the Breit (brick-wall) reference frame. In fact, to this order in the LEX the formula is valid for ω and θ being the energy and angle in the lab or center-of-mass frame.

Equation (6) shows that the leading (in LEX) effect of the electric polarizability cancels out, while the magnetic polarizability remains. Hence, our first claim is that a low-energy measurement of Σ_3 can in principle be used to extract β_{M1} independently of α_{E1} , just as it was proposed for the combination of polarised cross sections given in Eq. (4).

In reality the low-energy Compton experiments on the proton are difficult because of small cross sections and overwhelming QED backgrounds. Precision measurement only become feasible for photon-beam energies above 60 MeV and scattering angles greater than 40 degrees. The upcoming experiments at HIGS and MAMI are planned for the energies between 80 and 150 MeV. As mentioned above, at these energies the effect of higher-order terms may become substantial. One way to see it is to compare the LEX result with the dispersion-relation calculations or calculations based on chiral perturbation theory.

Figures 2 and 3 demonstrate such a comparison of the leading-LEX result to the next-next-to-leading order (NNLO) BChPT result of Ref. [4] for the two observables defined in Eqs. (4) and (5). The observables are plotted for the case of proton Compton scattering as a function of magnetic polarizability of the proton. From Fig. 2 one sees that for the beam energy of 100 MeV the LEX is in a good agreement with the BChPT result, especially for the forward directions (left panels).




FIG. 2: (Color online). Beam asymmetry Σ_3 — upper panel, and the linear combination of polarised cross sections defined in Eq. (4) — lower panel, shown as function of β_{M1} for fixed photon energy of 100 MeV and scattering angles of 60° (left panels) and 120° (right panels) degrees. The curves are: dashed green — Born contribution; dash-dotted magenta — the leading LEX formula Eq. (6); red solid — NNLO BChPT [4].

¹ The problem seems to be very similar to the case of proton form factors, where the angular (Rosenbluth) separation from unpolarized scattering did not yield the correct result for the electric form factor, due to the dominance of the magnetic contribution, and only separating the electric form factor from the magnetic one by using polarization has yielded a break through. See [16, 17] for reviews.




FIG. 3: (Color online). The same as in the previous figure but for photon beam energy of 135 MeV.

As expected we observe a significant sensitivity of these observables to β_{M1} . However, the conventional observable Eq. (4) is small compared to the unpolarised cross section (a few nano-barns per steradian compared to a few tens of nb/sr). It is therefore extremely difficult to measure it to the accuracy of few percent required here to discriminate between the PDG and ChPT values for the magnetic polarizability displayed in Fig. 1. In contrast, the beam asymmetry is large and, given the fact that many systematic errors tend to cancel out in this

observable, the required accuracy should be much easier to achieve.

The results for the beam energy of 135 MeV (Fig. 3) show that the leading LEX result is invalid at such energies. The differences between the LEX and BChPT curves at both 100 and 135 MeV are mainly due to α_{E1} (which here is equal to zero for the LEX and about 11 for the BChPT) and the π^0 -anomaly contribution. While it is well-known that the anomaly contribution is proportional to $t/(t-m_\pi^2)$ and hence is maximal at the backward angles, the α_{E1} contribution needs to be examined.

In order to uniquely define the ‘recoil corrections’ due to the polarizabilities, we ought introduce them in a Lorentz-invariant fashion by writing down an effective Lagrangian which yields the right Hamiltonian in the static limit, i.e.:

$$\begin{aligned} \mathcal{L}_{NN\gamma\gamma} = & \pi\beta_{M1}\bar{N}NF^2 \\ & - \frac{2\pi(\alpha_{E1} + \beta_{M1})}{M^2}(\partial_\alpha\bar{N})(\partial_\beta N)F^{\alpha\mu}F^{\beta\nu}\eta_{\mu\nu} \end{aligned} \quad (8)$$

where $F_{\mu\nu} = \partial_{[\mu}A_{\nu]}$ is the electromagnetic field-strength tensor, $N(x)$ is the nucleon Dirac-spinor field, $\eta_{\mu\nu} = \text{diag}(1, -1, -1, -1)$ is the Minkowski metric. Recalling that $F^2 = -2(F^{0i})^2 + (F^{ij})^2 = -2\mathbf{E}^2 + 2\mathbf{B}^2$, assuming the nucleon rest frame: $\partial_i N = 0$, $i\gamma^0\partial_0 N = MN$, $i\partial_0\bar{N}\gamma^0 = -M\bar{N}$, we obtain

$$\mathcal{L}_{NN\gamma\gamma} = 2\pi\{\beta_{M1}(\mathbf{B}^2 - \mathbf{E}^2) + (\alpha_{E1} + \beta_{M1})\mathbf{E}^2\}\bar{N}N,$$

which readily reproduces the well-known non-relativistic Hamiltonian: $4\pi(-\frac{1}{2}\alpha_{E1}\mathbf{E}^2 - \frac{1}{2}\beta_{M1}\mathbf{B}^2)$.

The Feynman amplitude corresponding to the Lagrangian in Eq. (8) can be written as:

$$\begin{aligned} \mathcal{M}_{\text{NB}}^{\mu\nu}\varepsilon'_\mu\varepsilon_\nu &= 4\pi\bar{u}(p')u(p)\left[\beta_{M1}(q\cdot q'\varepsilon'\cdot\varepsilon - q\cdot\varepsilon'q'\cdot\varepsilon) + \frac{\alpha_{E1} + \beta_{M1}}{2M^2}(p'_\alpha p_\beta + p_\alpha p'_\beta)(q'^\alpha\varepsilon'^\mu - q'^\mu\varepsilon'^\alpha)(q^\beta\varepsilon_\mu - q_\mu\varepsilon^\beta)\right] \\ &= \bar{u}(p')u(p)\left[-A_1^{(\text{NB})}(s,t)\mathcal{E}'\cdot\mathcal{E} + A_2^{(\text{NB})}(s,t)q\cdot\mathcal{E}'q'\cdot\mathcal{E}\right], \end{aligned} \quad (9)$$

where p and q (p' and q') are the four-momenta of incident (outgoing) nucleon and photon and the manifestly gauge-invariant polarization vectors are then:

$$\mathcal{E}_\mu = \varepsilon_\mu - \frac{(p' + p)\cdot\varepsilon}{(p' + p)\cdot q}q_\mu, \quad (10a)$$

$$\mathcal{E}'_\mu = \varepsilon'_\mu - \frac{(p' + p)\cdot\varepsilon'}{(p' + p)\cdot q}q'_\mu. \quad (10b)$$

The polarizability contribution to the invariant Compton amplitudes is thus given as follows:

$$A_1^{(\text{NB})}(s,t) = 2\pi(\alpha_{E1} + \beta_{M1})(\nu^2 + \nu'^2) + 2\pi\beta_{M1}t, \quad (11a)$$

$$A_2^{(\text{NB})}(s,t) = -4\pi\beta_{M1}. \quad (11b)$$

We note that the contribution of $\alpha_{E1} + \beta_{M1}$ to A_1 differs from conventional definitions by terms of higher order in the Mandelstam variable t , and hence in energy. For instance, the difference of the present $A_1^{(\text{NB})}$ with the one in Ref. [12]

is equal to: $-\pi(\alpha_{E1} + \beta_{M1})t(\omega^2 - \frac{1}{4}t)$. We however need to retain these terms to obtain the following LEX expression:

$$\begin{aligned} \Sigma_3 = & \Sigma_3^{(B)} - \frac{4 \cos \theta \sin^2 \theta}{(1 + \cos^2 \theta)^2} \frac{M^3 \beta_{M1}}{\alpha_{em}} \left(\frac{\omega}{M} \right)^2 \left\{ 1 + \left(\frac{\omega}{M} \right)^2 \left[a_1(\cos \theta) + \frac{M^3 \alpha_{E1}}{\alpha_{em}} \right] \right\} \\ & + \frac{\sin^2 \theta}{(1 + \cos^2 \theta)^2} \left(\frac{\omega}{M} \right)^4 \left[a_2(\cos \theta) \frac{M^3 \alpha_{E1}}{\alpha_{em}} + a_3(\cos \theta) \left(\frac{M^3 \beta_{M1}}{\alpha_{em}} \right)^2 \right] + O(\omega^6), \end{aligned} \quad (12)$$

where the dimensionless coefficient functions are given by (with κ denoting the anomalous magnetic moment):

$$\begin{aligned} a_1(z) &= -\frac{1}{4(1+z^2)} [7 - 8z + z^4 + 2(8 - 16z + 7z^2 - z^4)\kappa + (32 - 40z - z^2 - z^4)\kappa^2 \\ &\quad + 4(5 - 4z - 3z^2)\kappa^3 + (5 - 3z^2)\kappa^4], \\ a_2(z) &= 2(1 - z) + 2(2 - 4z + z^2)\kappa + (7 - 10z - 2z^2)\kappa^2 + 4(1 - z - z^2)\kappa^3 + (1 - z^2)\kappa^4, \\ a_3(z) &= \frac{2(1 - 6z^2 + z^4)}{1 + z^2}. \end{aligned} \quad (13)$$

At order ω^4 there are also contributions from higher polarizabilities; in the case of Σ_3 these are the spin polarizabilities and the magnetic quadrupole polarizability β_{M2} . Spin polarizabilities are dominated by the π^0 anomaly and hence their contribution is only appreciable in the backward angles. The contribution of β_{M2} is tiny below 100 MeV. As for the $O(\omega^4)$ contribution of scalar polarizabilities shown above, we note that the coefficient functions containing κ (i.e., a_1 and a_2) can become large in the backward angles. To limit the impact of these terms on the extraction of β_{M1} the scattering angle should be around 60 degrees, as is also seen in Fig. 2.

To conclude, the beam asymmetry Σ_3 is instrumental in isolating the contribution of the magnetic polarizability β_{M1} to low-energy Compton scattering. While the cross sections receive contributions from both the electric and magnetic polarizability, the effect of α_{E1} cancels out from the asymmetry at leading order in the low-energy expansion. We have shown that the next-to-leading corrections are minimal at the forward scattering angles. A precise and model-independent determination of the proton β_{M1} is feasible through a precision measurement of Σ_3 at beam energies of about 100 MeV and forward scattering angles. Furthermore, when multiplied with the unpolarized cross section, Σ_3 yields the polarized cross-section difference which provides an exclusive access to the electric polarizability. A measurement of the beam asymmetry at these low energies has not been done yet, but is being planned at the MAMI and HIGS facilities and could be done in the very near future.

Acknowledgements

We thank Dave Hornidge for helpful remarks on the manuscript. We acknowledge the support of the Deutsche

Forschungsgemeinschaft (DFG) through Collaborative Research Center SFB 1044 “The Low-Energy Frontier of the Standard Model” and Graduate School DFG/GRK 1581 “Symmetry Breaking in Fundamental Interactions”.

-
- [1] J. Beringer *et al.* [Particle Data Group Collaboration], Phys. Rev. D **86**, 010001 (2012).
 - [2] A. M. Baldin, Nucl. Phys. **18**, 310 (1960).
 - [3] B. R. Holstein, Comments Nucl. Part. Phys. **20**, 301 (1992).
 - [4] V. Lensky and V. Pascalutsa, Eur. Phys. J. C **65**, 195 (2010).
 - [5] J. A. McGovern, D. R. Phillips and H. W. Griesshammer, Eur. Phys. J. A **49**, 12 (2013).
 - [6] F. J. Federspiel *et al.*, Phys. Rev. Lett. **67**, 1511 (1991).
 - [7] A. Zieger, R. Van de Vyver, D. Christmann, A. De Graeve, C. Van den Abeele and B. Ziegler, Phys. Lett. B **278**, 34 (1992).
 - [8] B. E. MacGibbon, G. Garino, M. A. Lucas, A.M. Nathan, G. Feldman and B. Dolbilkin, Phys. Rev. C **52**, 2097 (1995).
 - [9] V. Olmos de Leon *et al.*, Eur. Phys. J. A **10**, 207 (2001).
 - [10] D. Babusci, G. Giordano and G. Matone, Phys. Rev. C **57**, 291 (1998).
 - [11] V. Lensky, J. A. McGovern, D. R. Phillips and V. Pascalutsa, Phys. Rev. C **86**, 048201 (2012).
 - [12] H. W. Griesshammer, J. A. McGovern, D. R. Phillips and G. Feldman, Prog. Part. Nucl. Phys. **67**, 841 (2012).
 - [13] J. L. Powell, Phys. Rev. **75**, 32 (1949).
 - [14] D. Drechsel, B. Pasquini and M. Vanderhaeghen, Phys. Rept. **378**, 99 (2003).
 - [15] M. Schumacher, Prog. Part. Nucl. Phys. **55**, 567 (2005).
 - [16] C. E. Hyde and K. de Jager, Ann. Rev. Nucl. Part. Sci. **54**, 217 (2004).
 - [17] C. F. Perdrisat, V. Punjabi, and M. Vanderhaeghen, Prog. Part. Nucl. Phys. **59**, 694 (2007).
 - [18] L. C. Maximon, Phys. Rev. C **39**, 347 (1989).